

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/00, 48/00		A1	(11) International Publication Number: WO 95/24185
			(43) International Publication Date: 14 September 1995 (14.09.95)
(21) International Application Number: PCT/US94/02652		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 11 March 1994 (11.03.94)		Published <i>With international search report.</i>	
(71) Applicant: ISIS PHARMACEUTICALS, INC. [US/US]; 2280 Faraday Avenue, Carlsbad, CA 92008 (US).			
(72) Inventors: ACEVEDO, Oscar; 12117 Brickellia, San Diego, CA 92129 (US). DUNKEL, Martin, G.; Rudolf-Wissel Strasse 28, D-37079 Gottingen (DE). COOK, Phillip, Dan; 7340 Bolero Street, Carlsbad, CA 92009 (US).			
(74) Agents: CALDWELL, John, W. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).			
(54) Title: NOVEL PYRIMIDINE NUCLEOSIDES			
(57) Abstract			
Novel 2- and 4-substituted pyrimidine nucleosides and oligonucleosides are provided, as are methods and intermediates useful in preparing same.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

WO 95/24185

PCT/US94/02652

NOVEL PYRIMIDINE NUCLEOSIDES

FIELD OF THE INVENTION

This invention relates to the design and synthesis of certain 2- and 4-substituted pyrimidine nucleosides that are
5 useful for incorporation into oligonucleotides. The oligonucleotides, in turn, are useful as diagnostic reagents, research reagents and in therapeutics.

BACKGROUND OF THE INVENTION

It is well known that most of the bodily states in
10 multicellular organisms, including most disease states, are effected by proteins. Such proteins, either acting directly or through their enzymatic or other functions, contribute in major proportion to many diseases and regulatory functions in animals and man. Classical therapeutics has generally focused upon
15 interactions with such proteins in efforts to moderate their disease-causing or disease-potentiating functions. In newer therapeutic approaches, modulation of the actual production of such proteins is desired. By interfering with the production of proteins, the maximum therapeutic effect might be obtained
20 with minimal side effects. It is the general object of such therapeutic approaches to interfere with or otherwise modulate

WO 95/24185

2

PCT/US94/02652

gene expression which would lead to undesired protein formation.

One method for inhibiting specific gene expression is with the use of oligonucleotides. Oligonucleotides complementary to a specific target messenger RNA (mRNA) sequence are used. Several oligonucleotides are currently undergoing clinical trials for such use.

Transcription factors interact with double-stranded DNA during regulation of transcription. Oligonucleotides can serve as competitive inhibitors of transcription factors to modulate the action of transcription factors. Several recent reports describe such interactions (see, Bielinska, et. al., *Science* 1990, 250, 997-1000; and Wu, et al., *Gene* 1990, 89, 203-209.)

Oligonucleotides also have found use in diagnostic tests. Such diagnostic tests can be performed using biological fluids, tissues, intact cells or isolated cellular components. As with the above gene expression inhibition, diagnostic use can take advantage of an oligonucleotides ability to hybridize with a complementary strand of nucleic acid. Hybridization is the sequence specific hydrogen bonding of oligonucleotides via Watson-Crick and/or Hoogsteen base pairs to RNA or DNA. The bases of such base pairs are said to be complementary to one another.

Oligonucleotides are also widely used as research reagents. They are useful for understanding the function of many other biological molecules as well as in the preparation of such other biological molecules. One particular use, the use of oligonucleotides as primers in the reactions associated with polymerase chain reaction (PCR), has been the cornerstone for the establishment of an ever expanding commercial business. The use of such PCR reactions has seemingly "exploded" as more and more use of this very important biological tool are practiced. The uses of PCR have extended into many areas in addition to those contemplated by its nobel laureate inventor. Examples of such new areas include forensics, paleontology, evolutionary studies and genetic counseling to name just a few. Primers are needed for each of these uses. Oligonucleotides, both natural and synthetic, serve as the primers.

WO 95/24185

PC 7/US94/02652

- 3 -

Oligonucleotides also are used in other laboratory procedures. A number of these uses are described in common laboratory manuals such as *Molecular Cloning, A Laboratory Manual*, Second Ed., J. Sambrook, et al., Eds., Cold Spring Harbor Laboratory Press, 1989; and *Current Protocols In Molecular Biology*, F. M. Ausubel, et. al., Eds., Current Publications, 1993. Such uses include Synthetic Oligonucleotide Probes, Screening Expression Libraries with Antibodies and Oligonucleotides, DNA Sequencing, In Vitro Amplification of DNA by the Polymerase Chain Reaction and Site-directed Mutagenesis of Cloned DNA from Book 2 of *Molecular Cloning, A Laboratory Manual*, *ibid.* and DNA-Protein Interactions and The Polymerase Chain Reaction from Vol. 2 of *Current Protocols In Molecular Biology*, *ibid.*.

To supply the users of oligonucleotides, many scientific journals now contain advertisements for either oligonucleotide precursors or for custom-synthesized oligonucleotides. This has become an important commercial use of oligonucleotides. Other oligonucleotides can be synthesized to have properties that are tailored for the desired use. Thus, a number of chemical modifications have been introduced into oligonucleotides to increase their usefulness in diagnostics, as research reagents, and as therapeutic entities. These modifications are designed to increase binding to a target strand, to assist in identification of the oligonucleotide or an oligonucleotide-target complex, to increase cell penetration, to provide stability against nucleases and other enzymes that degrade or interfere with the structure or activity of the oligonucleotides, to provide a mode of disruption (terminating event) once sequence-specifically bound to a target, or to improve the pharmacokinetic properties of the oligonucleotides.

To serve some of the above-noted uses, reactive functionalities can be tethered to an oligonucleotide and directed at phosphodiester and heterocyclic centers in target compounds. The positioning of these functionalities may require the synthesis of C-2 and C-4 substituted pyrimidine

WO 95/24185

PCT/US94/02652

- 4 -

nucleosides with the following general features: (a) preservation of Watson-Crick hybridization characteristics; (b) stability during automated DNA synthesis; and (c) the conformational rigidity of a carbon tether to accurately place
5 functionalities near their target. Criteria (a) and (b) are not met by readily available 2-O-, 2-S-, or 2-N-alkyl pyrimidine nucleosides since these groups are sensitive to nucleophilic displacement. For example, deprotecting DNA
10 oligomers containing S-alkylated moieties with concentrated ammonia likely would yield displacement products at C-2 and C-4 pyrimidine positions. In addition, 2-O-, 2-S-, and 2-N-alkyl groups would preclude the availability of an imino proton at N-3, which is necessary for hydrogen bonding. Likewise,
15 nucleosides having 2-N-alkyl groups at these positions would fail to meet criteria (b) and (c) due to their chemical instability and the spatial instability inherent in an inversion of configuration of a tertiary nitrogen. Accordingly, there remains a need in the art for novel 2- and
20 4-substituted pyrimidines that meet the design criteria set forth above.

OBJECTS OF THE INVENTION

It is one object of this invention to provide oligonucleotides for use in diagnostics, therapeutics and as
research reagents.

25 It is another object of this invention to provide oligonucleotides that effectively modulate the activity of RNA or DNA or proteins.

It is a further object to provide novel 2-substituted pyrimidines.

30 These and other objects will become apparent to persons of ordinary skill in the art from a review of the present specification and appended claims.

SUMMARY OF THE INVENTION



In accordance with these and other objects, the
35 present invention provides novel C-2 and C-4 modified

WO 95/24185

PCT/US94/02652

- 5 -

pyrimidine nucleosides and oligonucleosides. In certain embodiments, the compounds of the invention have structure 1 or structure 2:



wherein:

- 5 J is N or CH;
 R₅ is H or CH₃;
 one of R₂ and R₄ is =O, =NH, or =NH₂⁺; and the other of
 R₂ and R₄ is Q, =C(R_A)-Q, C(R_A)(R_B)-C(R_C)(R_D)-Q, C(R_A)=C(R_C)-Q or
 C≡C-Q;
- 10 R_A, R_B, R_C and R_D independently are H, SH, OH, NH₂, or
 C₁-C₂₀ alkyl, or one of (R_A)(R_B) or (R_C)(R_D) is =O;
 Q is halogen, hydrogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkylamine,
 C₁-C₂₀ alkyl-N-phthalimide, C₁-C₂₀ alkylimidazole, C₁-C₂₀
 alkylbis-imidazole, imidazole, bis-imidazole, amine, N-
 15 phthalimide, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, hydroxyl, thiol,
 keto, carboxyl, nitrates, nitro, nitroso, nitrile, trifluor-
 omethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-
 dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, azido, hydrazino,
 hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide,
 20 disulfide, silyl, O-(hydroxyl protecting group), a leaving
 group, a heterocycle, an intercalator, a reporter molecule, a
 conjugate, a polyamine, a polyamide, a polyethylene glycol, a
 polyether, a group that enhances the pharmacodynamic properties
 of oligonucleotides, a group that enhances the pharmacokinetic
 25 properties of oligonucleotides, a RNA cleaving moiety or a
 depurination enhancing group;

95/2418

PCT/US94/02652

- 6 -

T_1 and T_2 independently are H, phosphate, an activated phosphate, a hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleotide or an oligonucleoside;

X_1 and X_2 independently are hydrogen, halogen, hydroxyl, O-(hydroxyl protecting group), thiol, carboxyl, nitrate, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocyclic, alicyclic, carbocyclic, intercalators, reporter molecules, conjugates, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligonucleotides, or groups that enhance the pharmacokinetic properties of oligonucleotides.

For structure 1, when T_2 is a hydroxyl protecting group or together T_1 and one of X_1 or X_2 is a hydroxyl protecting group, then R_2 should not be CH_3 ; when T_1 and T_2 are OH or O-benzoyl, then R_2 should not be $CH_2OC_6H_5$; and when R_4 is Q then Q is not hydrogen, hydroxyl, thiol, amino or hydrogen.

The present invention also provides methods for preparing the compounds of the invention and synthetic intermediates employed in the methods. In certain embodiments, the methods involve reacting 2,2'- or 2,5'-anhydropyrimidines with a nucleophile to effect attack of the nucleophile at the 2-pyrimidine position. The nucleophile can have general formula R_2 or can be a functional group that can be modified to have formula R_2 . In further embodiments, the methods involve reacting 4-halopyrimidines with substituted alkynes in the presence of an organometallic coupling reagent for a time and under reaction conditions effective to form a covalent bond between the alkyne and the pyrimidine 4-position.

The invention also provides methods for modulating the production or activity of a protein in an organism comprising contacting the organism with a compound according to the invention. It is preferred that the compound include a base sequence that is complementary to a RNA or DNA sequence that codes for the protein. This "targeting portion" of the

WO 95/24185

PCT/US94/02652

- 7 -

compound is, thus selected to be complementary to the targeted portion of DNA or RNA such that the compound can bind to the target portion.

BRIEF DESCRIPTION OF THE DRAWING

5 Fig. 1 shows certain synthetic methods according to the present invention.

DETAILED DESCRIPTION OF INVENTION

In a general sense, the present invention is directed to nucleosides that include 2- and 4-modified pyrimidine bases, and to more complex compounds that include the nucleosides. The term "nucleoside" as used in connection with this invention refers to a unit made up of a heterocyclic base and its sugar. The term "nucleotide" refers to a nucleoside having a phosphate group on its 3' or 5' sugar hydroxyl group. Thus, nucleosides, unlike nucleotides, have no phosphate group. "Oligonucleotide" refers to a plurality of joined nucleotide units formed in a specific sequence from naturally occurring bases and pentofuranosyl groups joined through a sugar group by native phosphodiester bonds. This term refers to both naturally occurring and synthetic species formed from naturally occurring subunits.

The compounds of the invention generally can be viewed as "oligonucleotide analogs", that is, compounds which function like oligonucleotides but which have non-naturally occurring portions. Oligonucleotide analogs can have altered sugar moieties, altered base moieties or altered inter-sugar linkages. For the purposes of this invention, an oligonucleotide analog having non-phosphodiester bonds, i.e., an altered inter-sugar linkage, is considered to be an "oligonucleoside." The term "oligonucleoside" thus refers to a plurality of nucleoside units joined by linking groups other than native phosphodiester linking groups. The term "oligomers" is intended to encompass oligonucleotides, oligonucleotide analogs or oligonucleosides. Thus, in speaking of "oligomers" reference is made to a series of nucleosides or

WO 95/24184

PCT/US94/02652

- 8 -

nucleoside analogs that are joined via either naturally occurring phosphodiester bonds or other, non-naturally occurring linkages.

The oligonucleosides of the invention preferably comprise from about 5 to about 50 nucleosides, each nucleoside having the same or different structure. The targeting portions of the oligonucleosides preferably comprise about 8 to about 40 base nucleosides, more preferably about 12 to 20 base nucleosides.

It is preferred that an oligonucleoside of the invention include at least one 2- or 4-modified pyrimidine at its 3'- or 5'-terminus. In certain embodiments, the oligonucleoside has 1-3 modified pyrimidines at its 3'-terminus.

The nucleosides of the invention can be linked with one another or with other nucleosides or oligonucleosides via a sugar linking group. The linking groups can be any structure capable of linking sugar moieties of oligonucleosides to form the targeting portion of the compositions of this invention.



It is preferred that these sugar linking groups comprise phosphodiester groups or derivatives thereof. Derivatives of the phosphodiester structure can include substitution of a sulphur, methyl, methyl oxide, or amine group for an oxygen. The sugar phosphate nucleic acid backbone may be modified as a phosphorothioate, methylphosphonate, or phosphate alkylated moiety. The phosphodiester linkage may also be replaced by a carbon or ether linkage.

In certain embodiments, the 2-substituted pyrimidine compounds of the invention are represented by the structures 1 and 2:

WO 95/24185

PCT/US94/02652

- 9 -



wherein:

J is N or CH;

R₅ is H or CH₃;

one of R₂ and R₄ is =O, =NH, or =NH₂⁺; and the other of

5 R₂ and R₄ is Q, =C(R_A)-Q, C(R_A)(R_B)-C(R_C)(R_D)-Q, C(R_A)=C(R_C)-Q or C≡C-Q;

R_A, R_B, R_C and R_D independently are H, SH, OH, NH₂, or C₁-C₂₀ alkyl, or one of (R_A)(R_B) or (R_C)(R_D) is =O;

Q is halogen, hydrogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkylamine, 10 C₁-C₂₀ alkyl-N-phthalimide, C₁-C₂₀ alkylimidazole, C₁-C₂₀ alkylbis-imidazole, imidazole, bis-imidazole, amine, N-phthalimide, C₂-C₂₀ alkenyl, C₂-C₂₀ alkynyl, hydroxyl, thiol, keto, carboxyl, nitrates, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N- 15 dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, O-(hydroxyl protecting group), a leaving group, a heterocycle, an intercalator, a reporter molecule, a conjugate, a polyamine, a polyamide, a polyethylene glycol, a 20 polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, a group that enhances the pharmacokinetic properties of oligonucleotides, a RNA cleaving moiety or a depurination enhancing group;

T₃ and T₅ independently are H, phosphate, an activated 25 phosphate, a hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleotide or an oligonucleoside;

WO 95/24183

PCT/US94/02652

- 10 -

X_1 and X_2 independently are hydrogen, halogen, hydroxyl, O-(hydroxyl protecting group), thiol, carboxyl, nitrate, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocyclic, alicyclic, carbocyclic, intercalators, reporter molecules, conjugates, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligonucleotides, or groups that enhance the pharmacokinetic properties of oligonucleotides.

For structure 1, when T_5 is a hydroxyl protecting group or together T_3 and one of X_1 or X_2 is a hydroxyl protecting group, then R_2 should not be CH_3 ; when T_3 and T_5 are OH or O-benzoyl, then R_2 should not be $CH_2OC_6H_5$.

Preferred R_2 and R_4 groups include $C(R_A)=C(R_C)-Q$, preferably where R_A is OH, SH or NH_2 and R_C is H, more preferably where Q is C_{1-20} or C_{1-6} alkyl; and $C(R_A)(R_B)-C(R_C)(R_D)-Q$, preferably where R_A and R_B are both hydrogen and R_C and R_D together are =O.

Preferred structures are also those in which one of X_1 or X_2 is hydrogen and the other is hydroxyl, or where both are hydrogen; where R_4 is =O; where R_5 is hydrogen or CH_3 ; or where T_3 is an activated phosphate and T_5 is a hydroxyl protecting group; or where one of T_3 or T_5 is an oligonucleotide or an oligonucleoside, or both are H.

A wide variety of protecting groups can be employed in the methods of the invention. See, e.g., Beaucage, et al., Tetrahedron 1992, 12, 2223. In general, protecting groups render chemical functionality inert to specific reaction conditions, and can be appended to and removed from such functionality in a molecule without substantially damaging the remainder of the molecule. Representative hydroxyl protecting groups include t-butyldimethylsilyl (TBDMS), t-butyldiphenylsilyl (TBDPS), dimethoxytrityl (DMTr), monomethoxytrityl (MMTr), and other hydroxyl protecting groups as outlined in the above-noted Beaucage reference.

WO 95/24185

PCT/US94/02652

- 11 -

Leaving groups according to the invention are chemical functional groups that can be displaced from carbon atoms by nucleophilic substitution. Representative leaving groups include but are not limited to halogen, alkylsulfonyl, substituted alkylsulfonyl, arylsulfonyl, substituted arylsulfonyl, heterocyclosulfonyl or trichloroacetimidate groups. Preferred leaving groups include chloro, fluoro, bromo, iodo, p-(2,4-dinitroanilino)benzenesulfonyl, benzenesulfonyl, methylsulfonyl (mesylate), p-methylbenzenesulfonyl (tosylate), p-bromobenzenesulfonyl, trifluoromethylsulfonyl (triflate), trichloroacetimidate, acyloxy, 2,2,2-trifluoroethanesulfonyl, imidazolesulfonyl, and 2,4,6-trichlorophenyl groups.

Heterocycles according to the invention are functional groups that include atoms other than carbon in their cyclic backbone.

Intercalators according to the invention generally include non-carcinogenic, polycyclic aromatic hydrocarbons or heterocyclic moieties capable of intercalating between base pairs formed by a hybrid oligonucleotide/RNA target sequence duplex. Intercalators can include naphthalene, anthracene, phenanthrene, benzonaphthalene, fluorene, carbazole, acridine, pyrene, anthraquinone, quinoline, phenylquinoline, xanthene or 2,7-diazaanthracene groups. Other intercalators believed to be useful are described by Denny, *Anti-Cancer Drug Design* 1989, 4, 241. Another intercalator is the ligand 6-[[[9-[[6-(4-nitrobenzamido)hexyl]amino]acridin-4-yl]carbonyl]-amino]hexanoyl-pentafluorophenyl ester.

Reporter molecules are those compounds that have physical or chemical properties that allow them to be identified in gels, fluids, whole cellular systems, broken cellular systems and the like utilizing physical properties such as spectroscopy, radioactivity, colorimetric assays, fluorescence, and specific binding. Particularly useful reporter molecules include biotin and fluorescein dyes. Particularly useful as reporter molecules are biotin, fluorescein dyes, alkaline phosphates, and horseradish peroxidase.

WO 95/24185

PCT/US94/02652

- 12 -

The term "depurination enhancing moiety" includes chemical moieties that are capable of enhancing the rate of depurination of a purine-containing nucleic acid species. Depurination enhancing moieties enhance the rate of removal, break down, and/or loss of adenine and guanine nucleobases from adenosine and guanosine nucleotides. They also enhance the rate of the removal, break down, and/or loss of other purine-containing nucleotides such as 7-methylguanosine, 3-methylguanosine, wyosine, inosine, 2-aminoadenosine, and other "minor" or synthetic nucleotides. Preferred depurination enhancing moieties are sulfur-containing compounds, including sulfur-containing heterocycles and both cyclic and alicyclic sulfonium compounds. Specific examples include but are not limited to thiophene, thianthrene, isothiazole, alkyl sulfonium salts, thiophenium salts, 1,3-thiazolium salts, 1,2-oxathiolanium salts, alkyl 1,4-dithianium salts, alkyl thiazolium salts, thioniabicyclo[2,2,1]heptane salts and 3aH-1,6-dithia-6a-thioniapentalene salts. Anions for such salts include halide anions and other anions.

Conjugates are functional groups that improve the uptake of the compounds of the invention. Representative conjugates include steroid molecules, reporter molecules, non-aromatic lipophilic molecules, reporter enzymes, peptides, proteins, water soluble vitamins, and lipid soluble vitamins, as disclosed by U.S. Patent Application Serial No. 782,374, filed October 24, 1991, and PCT Application US92/09196, filed October 23, 1992, the disclosures of which are incorporated herein by reference. Representative conjugates also are disclosed by Goodchild, *Bioconjugate Chemistry* 1990, 1, 165.


Activated phosphates according to the invention are disclosed by Beaucage, *et al.*, *Tetrahedron* 1992, 2223.

As will be recognized by those of ordinary skill in the art, the base portion of structure 1 can be equivalently represented by its tautomeric forms. Thus, the recitation of structure 1 in connection with the present invention is intended to include each of the following tautomeric structures:

WO 95/4185

PCT/US94/02652

- 13 -



There appear to have been few reported syntheses of 2-alkyl substituted pyrimidine nucleosides. Witkop, et al., *J. Amer. Chem. Soc.* 1969, 91, 7751 reported that treatment of a protected 2,5'-anhydrouridine with dimethylsulfoxonium methylid (DMSY) gave a 2-methylpyrimidin-4-one riboside after hydro-

5 genolysis of the sulfoxonium group. 2-Methyl-pyrimidin-4-one ribonucleosides were also obtained when Rosenthal, et al., *Carbohydr. Res.* 1980, 78, 33, examined the reaction of 2,2'-anhydrouridines with 1,3-dithiane/*n*-butyllithium followed by

10 desulfurization. Vorbrüggen, et al., *Angew. Chem. Int. Ed.* 1976, 15, 689, have investigated the Eschenmoser sulfide extraction of 2- and 4-S-phenacylated-2-thiouridines to yield 2- and 4-phenacetylpyrimidin-4-one ribonucleosides. However, these methods presented a number of problems. For example,


15 attempts to synthesize the target nucleosides by DMSY chemistry according to Witkop's work resulted in laborious purifications and low yields. The Eschenmoser sulfide contraction is limited by the stability and limited availability of functionalized α -haloketones.

20 In certain embodiments, 2-substituted pyrimidine nucleosides of the invention are synthesized by nucleophilic substitution of 2,2'- or 2,5'-anhydropyrimidines. One preferred reaction sequence is shown in Scheme 1.

WO 93/24185

PCT/US94/02652

- 14 -



Scheme 1

At room temperature, the reaction of the lithium enolates of acetone, methylethylketone, 2-pentanone or 2-hexanone with a 3'-O-t-butyldiphenylsilyl-2,5'-anhydrothymidine 5 (T_3 = TBDMS, X_1 = H, R_5 = CH_3 or H) yields the corresponding 2-(β -ketoalkyl)-pyrimidin-4-one deoxynucleosides 6a-d (X_1 = H; T_5 = H; R_2 = $=\text{CHC}(\text{O})\text{CH}_3$ (6a), $=\text{CHC}(\text{O})\text{CH}_2\text{CH}_3$ (6b), $=\text{CHC}(\text{O})(\text{CH}_2)_2\text{CH}_3$ (6c), $=\text{CHC}(\text{O})(\text{CH}_2)_3\text{CH}_3$ (6d)) following removal of the protecting groups. Likewise, reaction of the lithium enolate of hexanone with a protected 2-5'-anhydrouridine 7 (T_3 = TBDMS, X_1 = OTBDMS, R_5 = CH_3 or H) yields the corresponding 2-(2-hexanon-1-yl)pyrimidin-4-one ribonucleoside 8 (X_1 = ribo OH, T_5 = T_3 = H, R_5 = CH_3 or H, R_2 = $=\text{CHC}(\text{O})(\text{CH}_2)_3\text{CH}_3$). The nucleoside structures 6a-d and 8 were isolated as yellowish oils with characteristic ultraviolet maxima near 320 nm; nucleoside structures 6a-d were found to be unstable. The downfield shift of the amidine/enol signal in the ^1H -NMR spectra of these compounds is indicative of protons bound by N-3 and keto oxygen. Further, two tautomeric structures can be drawn which account for the vinylic character of the alpha proton of the tether. 3',5'-(di-O-MMTr, -DMTr, -TBDMS or -TBDPS)-2,2'-anhydrouridines 9 (T_3 = T_5 = MMTr, DMTr, TBDMS, TBDPS) can be used to synthesize the corresponding arabinosides. The reaction of structure 9 (T_3 = T_5 = TBDMS) with the lithium enolate of 2-hexanone followed by a deprotection of the intermediate produced the 2-(2-hexanon-1-

WO 95/24185

PCT/US94/02652

- 15 -

yl)pyrimidin-4-one arabinonucleoside structure 10 (X_1 = ara OH, $T_2 = T_3 = H$, $R_5 = CH_3$ or OH, $R_2 = =CHC(O)(CH_2)_3CH_3$) in good yield. Also, it has been found that reaction of structure 9 ($T_2 = T_3 = TBDMS$) with the lithium enolate of 2-pentanone under standard conditions yields the protected 2-alkylated arabinonucleoside 11 (X_1 = ara OH, $T_2 = T_3 = H$, $R_5 = CH_3$ or H, $R_2 = =CHC(O)(CH_2)_2CH_3$) in 75% yield.

The 4-substituted pyrimidine compounds of the invention are synthesized by metal-mediated coupling 2-halopyrimidines and substituted alkynes. One preferred procedure is shown in Figure 1. In this approach, terminal alkynes are coupled to a 4-chloropyrimidine ribonucleoside by methods generally in accordance with Heck, "Palladium Reagents in Organic Synthesis" Academic Press, Orlando, FL (1985). A 4-chloro-2-trimethylsilylethoxy-pyrimidine, structure 12, can be prepared by reaction of trimethylsilylethanol/n-butyllithium and 2,4-dichloropyrimidine in tetrahydrofuran (THF) at $-20^\circ C$. Condensation of structure 12 and 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose is carried out using TMS-triflate/ CH_3CN generally according to the procedures of Vorbruggen, et al., Chem. Ber. 1981, 114, 1234. Coupling of the 4-chloro-1-(2,3,5-tri-O-benzoyl- β -D-ribofuranosyl)pyrimidin-2-one, structure 13, with an alkyne, such as 1-hexyne, and catalyzed by bis-triphenylphosphine palladium (II) dichloride/copper (I) iodide provides 4-(1-hexyn-1-yl)pyrimidin-2-one ribonucleoside, structure 14a ($R = C_4H_9$). Similarly, the reaction of structure 13 with trimethylsilylacetylene provides 4-trimethylsilylethynylpyrimidin-2-one ribonucleoside, structure 14b ($R = TMS$).

Conjugated alkyne nucleoside structures 14a,b can be reacted with H_2S to yield 4-(2-thiol-1-alken-1-yl)pyrimidin-2-one ribonucleoside structures 16a ($R = C_4H_9$) and 16b ($R = TMS$), or with methanolic ammonia to yield deprotected 4-(2-amino-1-alken-1-yl)pyrimidin-2-one riboside structures 17a ($R = C_4H_9$) and 17b ($R = TMS$). Interestingly, the reaction of the 4-(2-thiol-1-hexen-1-yl)pyrimidin-2-one, 16, with ammonia displaces the thiol group and produces the enamine structure 17a.

WO 95/14:85

PCT/US94/02652

- 16 -

identical to the product obtained from the reaction of alkyne 14a with ammonia. Hydration of structure 14a followed by deprotection of intermediate structure 15a (R_1 = benzoyl) yields the enolized 4-(2-hydroxy-1-hexen-1-yl)pyrimidin-2-one
5 ribonucleoside, structure 18a (R_1 = H).

The enol nature of the 2-(β -ketoalkyl)- and 4-(β -ketoalkyl)pyrimidinone nucleoside analogs (6a-d, 8, 10, 11 and 18a) offer the potential for Watson-Crick hybridization. Because of this potential they can be considered as analogues
10 of thymidine and cytidine. Thus hybridization of these compounds to adenosine and guanosine in a bifurcated bonding scheme is possible.

The compounds of this invention can be used in diagnostics, therapeutics, and as research reagents and kits.
15 In certain uses in diagnostics and as research reagents, the pyrimidine nucleosides of the invention are converted to their corresponding phosphoramidate nucleotide and incorporated into oligonucleotides. For incorporation into oligonucleotides, they can be used in place of a standard or normal nucleotide to
20 alter the properties of the diagnostic oligonucleotide reagent, as for instance, to increase the nuclease resistances of the oligonucleotide or to modulate (i.e., increase or decrease) hybridization of the oligonucleotide to a target.

Oligonucleotides that incorporate the compounds of the
25 invention can be used for the selective binding of RNA for other research and diagnostic purposes. Such selective, strong binding is accomplished through interaction of RNA or DNA with compounds of the invention that are resistant to degradative nucleases and which hybridize more strongly and with greater
30 fidelity than known oligonucleotides or oligonucleotide analogs.

For certain uses of the compounds of the invention, be it diagnostic use or therapeutic use, it is preferred that a target RNA or DNA be preselected to comprise that portion of
35 DNA or RNA which codes for a protein whose formation or activity is to be modulated. The targeting portion of the

WO 95/24185

PCT/US94 02652

- 17 -

compound to be employed is, thus, selected to be complementary to the targeted portion of DNA or RNA.

In accordance with a further preferred embodiment, the compounds of the invention mimic the secondary structure of natural nucleic acid structures that bind to particular proteins. The nucleic acid mimic can then be used for isolation and identification of the protein structure or for competitive binding to the protein structure. One such nucleic acid structure is the TAR structure, which binds the tat protein. By effectively mimicking TAR, compounds according to the invention bind the tat protein and thereby form a complex that can be isolated and identified using affinity chromatography.

For therapeutic use, oligonucleotide analogs are administered to animal suffering from a disease modulated by some protein. It is preferred to administer to patients suspected of suffering from such a disease an amount of oligonucleotide analog that is effective to reduce the symptomology of that disease. One skilled in the art can determine optimum dosages and treatment schedules for such treatment regimens.

In accordance with preferred embodiments, compounds of the invention hybridize to the mRNA of various pathogenic organisms. Preferred compounds for such use include those incorporating a complementary sequences for herpes, papilloma and other viruses.

It is generally preferred to administer the therapeutic agents in accordance with this invention internally such as orally, intravenously, or intramuscularly. Other forms of administration, such as transdermally, topically, or intra-lesionally may also be useful. Inclusion in suppositories may also be useful. Use of pharmacologically acceptable carriers is also preferred for some embodiments.

Additional objects, advantages, and novel features of this invention will become apparent to those skilled in the art upon examination of the following examples thereof, which are

WO 95/24185

PCT/US94/1262

- 18 -

not intended to be limiting, wherein parts and percents are by weight unless otherwise indicated.

Example 1

4-Chloro-2-trimethylsilylethoxy-pyrimidine, 12.

5 A solution of trimethylsilylethanol (10 g, 85 mmol) in 30 mL anhydrous THF was cooled to -68°C in a dry ice/isopropanol bath. A solution of n-butyl lithium (33.8 mL, 2.5 M in THF) was added dropwise over 15 min. The solution was stirred for 10 min at this temperature and then added to a
10 solution of 2,4-dichloropyrimidine (12.62 g, 85 mmol) in 75 mL THF at -35 to -25°C . The final yellowish solution was allowed to come to room temperature under an argon atmosphere then stirred at this temperature for 1 hr. This solution was diluted with 200 mL of diethylether and washed with cold water
15 (100 mL), saturated NaHCO_3 (100 mL) and brine, then dried over MgSO_4 , filtered and evaporated to yield a tan syrup. This syrup may be crystallized from hexanes at -60°C to yield 12 as a light yellow solid, 17.8 g (91 %). $^1\text{H-NMR}$ (CDCl_3): δ 8.36 (d, 1 H, H-6); 6.93 (d, 1 H, 5-H); 4.43 and 1.17 (t,t; 4 H; CH_2CH_3);
20 1.06 (bs, 9 H. TMS).

Example 2

4-Chloro-1-(2,3,5-tri-O-benzoyl- β -D-ribofuranosyl)pyrimidin-2-one, 13.

A solution of 12 (8.17 g, 35.4 mmol) in 200 mL
25 anhydrous acetonitrile is treated with 1-O-acetyl-2,3,5-tri-O-benzoyl-D-ribofuranose (Pfanstiehl) at room temperature and under an atmosphere of argon. To this mixture is added TMS-triflate (6.11 mL, 35.4 mmol) and the mixture stirred at this temperature for 2 hr. Evaporation of the solvents in vacuo
30 yielded an amber syrup which was immediately redissolved in 300 mL CH_2Cl_2 , washed with cold, saturated NaHCO_3 (2 x 150 mL), dried over magnesium sulfate, filtered and evaporated to a tan residue. Trituration of this residue with cold, anhydrous acetonitrile yielded analytically pure 13, 14.4 g (71 %). $^1\text{H-NMR}$ (DMSO-d_6): δ 8.39 (d, 1 H, 6-H, $J_{6,5}=11.8\text{ Hz}$); 8.0-7.4 (m,
35

WO 95/24185

PCI/US94/02652

- 19 -

15 H, aromatic); 6.68 (d, 1 H, H-5); 6.11 (d, 1H, H-1', $J_{1,2}=2$ Hz); 5.95 (m, 2 H, H-2', 3'); 4.85 (m, 1 H, H-4'); 4.70 (m, 2 H, H-5', 5"). Anal. Calcd for $C_{16}H_{23}N_2O_8Cl$: C, 62.67; H, 4.03; N, 4.87. Found: C, 62.21; H, 3.92; N, 4.80.

5 Example 3

4-(1-Hexyn-1-yl)-1-(2,3,5-tri-O-benzoyl- β -D-ribofuranosyl)-pyrimidin-2-one, 14a.

A suspension of bis-triphenylphosphine palladium (II) dichloride (0.35 g, 0.5 mmol) and copper (I) iodide (95 mg, 0.5 mmol) in THF (45 mL, distilled from Na-benzophenone ketyl) is sparged with dry argon gas for 10 min. To this suspension is added 1-hexyne (2.46 g, 30 mmol) followed by triethylamine (5 mL). The suspension is stirred at room temperature under argon for 15 min. A similarly sparged solution of the 2-chloro-1-(2,3,5-tri-O-benzoyl- β -D-ribofuranosyl)pyrimidin-4-one (13, 5.75 g, 10 mmol) in THF (15 mL) is added to the suspension of catalyst and the mixture stirred for 1 hr. The dark mixture is filtered through Celite and the filtrate evaporated in vacuo to afford a dark syrup. This syrup was triturated with cold diethylether, filtered and then dissolved in dichloromethane. The solution is washed with cold 1 N HCl, dried over magnesium sulfate and again evaporated in vacuo. The resulting residue was triturated with methanol to yield 14a as a white solid, 5.5 g, 88 %. 1H -NMR (DMSO- d_6): δ 8.33 (d, 1 H, H-6, $J_{6,5}=7$ Hz); 8.1-7.4 (m, 15 H, aromatic); 6.53 (d, 1 H, H-5); 6.24 (s, 1 H, H-1'); 2.52, 1.50 and 0.94 (3 m, 9 H, n-butyl). Anal. Calcd for $C_{36}H_{32}N_2O_8$: C, 69.67; H, 5.19; N, 4.51. Found: C, 69.32; H, 5.13; N, 4.44.

Example 4

30 2-(2-Trimethylsilylethynyl)-1-(2,3,5-tri-O-benzoyl- β -D-ribofuranosyl)pyrimidin-2-one, 14b.

Coupling of chloro nucleoside 13 (5.74 g, 10 mmol) with trimethylsilyl acetylene (4.9g, 50 mmol) according to the procedure described above gave 14b as a white amorphous solid, 35 5.12 g (80 %). 1H -NMR (DMSO- d_6): 8.37 (d, 1 H, H-6, $J_{6,5} = 6.5$

WO 95/24185

PCT/US94/02652

- 20 -

Hz); 8.1-7.4 (m, 15 H, aromatic); 6.60 (d, 1 H, H-5); 6.22 (s, 1 H, H-1'); 0.29 (s, 9 H, TMS). Anal. Calcd for $C_{32}H_{32}N_2O_8Si$: C, 66.02; H, 5.06; N, 4.40. Found: C, 65.68; H, 4.93; N, 4.39.

5 Example 5

4-(2-Hexanon-1-yl)-1-(2,3,5-tri-O-benzoyl- β -D-ribofuranosyl)-pyrimidin-2-one, 15a.

A solution of 14a (0.77 g, 1.2 mmol) in 50 mL acetone is treated with a 0.1 M solution of $Hg(OAc)_2$ (15 mL) in aqueous acetic- H_2O (1:1) and the mixture stirred at room temperature for 18 hr. The reaction is further diluted with dichloromethane (100 mL), washed with cold saturated $NaHCO_3$ solution, dried over magnesium sulfate and evaporated in vacuo. The resulting residue is flash-chromatographed using chloroform/acetone 19:1 to afford a yellowish solid. This solid is triturated with cold methanol to afford 15a, 0.47 g, 59%. 1H -NMR (DMSO- d_6): 12.5 (s, 1 H, enol); 8.0-7.4 (m, 16 H, H-6 plus aromatic); 6.15 (d, 1 H, H-1', $J = 2$ Hz); 5.90 (m, 3 H, H-5, 2', 3'); 5.39 (s, 1 H, vinylic, $J_{vinylic, allylic} \approx <1$ Hz); 2.30, 1.47, 1.24 and 0.86 (4 m, 9 H, n-butyl). Anal. Calcd for $C_{36}H_{34}N_2O_9$: C, 67.70; H, 5.37; N, 4.39. Found: C, 67.51; H, 5.28; N, 4.28.

Example 6

4-(2-Thiol-1-hexen-1-yl)-1-(2,3,5-tri-O-benzoyl- β -D-ribofuranosyl)pyrimidin-2-one, 16a.

A solution of compound 14a (0.64 g) in THF is sparged with a steady stream of H_2S gas for 3 min. The solution is then evaporated in vacuo and flash chromatographed using $CHCl_3$ /ethyl acetate (9:1) to afford 16a as an amber foam, 0.25 g (37%). 1H -NMR (DMSO- d_6): δ 15.3 (s, 1H, thiol); 6.43 (s, 1 H, vinylic, $J_{vinylic, allylic} < 1$ Hz); 6.23 (m, 2 H, H-1' and H-5); 2.7, 1.8, 1.3 and 0.9 (4 m, 9 H, n-butyl). Anal. Calcd for $C_{36}H_{34}N_2O_8S$: C, 66.04; H, 5.23; N, 4.27; S, 4.89. Found: C, 66.10; H, 5.18; N, 4.13; S, 4.43.

WO 95/24185

PCT/US94/02652

- 21 -

Example 7

4-(2-Amino-1-hexen-1-yl)-1-β-D-ribofuranosyl-pyrimidin-2-one,
17a.

A solution of 14a (1.2 g, 1.9 mmol) in methanolic
5 ammonia (saturated at -20° C) was stirred in a pressure bottle
for 18 hr. Evaporation of the solvents and flash
chromatography of the residue using ethyl acetate/methanol
(4:1) gave 17 a as a light yellow solid, 0.47 g (77 %). ¹H-NMR
(DMSO-d₆): δ 10.2 and 7.7 (2 bs, 2 H, NH₂); 7.83 (d, 1 H, H-6,
10 J_{6,5} = 6.9 Hz); 5.78 (s, 1 H, H-1'); 4.72 (s, 1 H, vinylic);
2.17, 1.50, 1.30 and 0.90 (4 m, 9 H, n-butyl).

Example 8

4-(2-Amino-1-ethen-1-yl)-1-β-D-ribofuranosyl-pyrimidin-2-one,
17b

15 A solution of 14b (1.2 g, 1.88 mmol) in 30 mL
methanolic ammonia (saturated at -20° C) was stirred at room
temperature for 18 hr. The solvents were evaporated in vacuo
to a bright red syrup. This syrup was triturated with
diethylether (Et₂O) (3 x 30 mL) then redissolved in methanol
20 (MeOH) and reevaporated to give a solid orange foam. A portion
of this foam was chromatographed on two preparative thin layer
chromatography (TLC) plates using ethyl acetate/MeOH, 4:1 to
give a sample for NMR. The purity of this sample was not
sufficient to fit elemental analysis. ¹H-NMR (DMSO-d₆): δ 9.5
25 and 7.7 (2 bs, 2 H, NH₂); 7.92 (d, 1 H, H-6, J_{6,5} = 7 Hz); 7.0
(m, d w/D₂O exch., 1 H, H-(b-vinylic), J_{b-a} = 7.4 Hz); 5.77 (s,
1 H, H-1'); 5.98 (d, 1 H, H-5); 4.74 (d, 1 H, H-(a-vinylic).

Example 9

2-Thio-1-(3',5'-di-O-benzoyl-2'-O-methyl-β-D-ribofuranosyl)-
30 pyrimidin-4-one

2-Thiouracil (1.282 g, 10 mmol), a few crystals of
ammonium sulfate, a few drops of trimethylsilyl chloride and
1,1,1,3,3,3-hexamethyldisilazane (HMDS, 20 ml) were refluxed
overnight. The clear, greenish solution was evaporated under
35 the exclusion of moisture. A solution of 2-O-methyl-1,3,5-tri-

WO 95/24185

PCT/US94/02632

- 22 -

O-benzoyl- α -D-ribose (4.76 g, 10 mmol) in acetonitrile (50 ml) was added to give a clear solution. SnCl₄ is added and the mixture became turbid and then immediately became clear thereafter. The mixture was stirred for three hours and then
5 poured into a stirred mixture of saturated NaHCO₃ and dichloromethane (200/400 ml). The organic layer was separated, dried over Na₂SO₄, filtered and evaporated. The amorphous residue was crystallized from hot ethanol to yield colorless
10 needles (1.895 g, 39%). ¹H-NMR (CDCl₃): δ 10.3 (s, 1 H, NH); 7.8 (d, 1 H, H6); 6.7 (s, 1 H, H1'); 5.7 (d, 1 H, H5); 3.6 (s, 3 H, OCH₃).

Example 10

2-Thio-S-(aceton-1-yl)-1-(3',5'-di-O-benzoyl-2'-O-methyl- β -D-ribofuranosyl)-pyrimidin-4-one

15 Chloroacetone (0.35 ml, 4.35 mmol) was added to a solution of 2-thio-1-(3',5'-di-O-benzoyl-2'-O-methyl- β -D-ribofuranosyl)-pyrimidin-4-one in dichloromethane (920 ml) and triethylamine (0.81 ml, 5.8 mmol). After 16 hrs water was added and the organic layer washed, dried with Na₂SO₄, filtered
20 and evaporated to yield the product as a homogeneous material by TLC. ¹H-NMR (CDCl₃): δ 7.5 (d, 1 H, H6); 6.0 (s, 1 H, H1'); 5.8 (d, 1 H, H5); 4.05 (s, 1 H, CH₂); 3.4 (s, 3 H, OCH₃); 2.3 (s, 3 H, COCH₃).

Example 11

25 2-(Aceton-1-yl)-1-(3',5'-di-O-benzoyl-2'-O-methyl)- β -D-ribofuranosyl)pyrimidin-4-one

Crude 2-thio-S-(aceton-1-yl)-1-(3',5'-di-O-benzoyl-2'-O-methyl- β -D-ribofuranosyl)-pyrimidin-4-one was dissolved in toluene (20 ml) and triphenylphosphine and potassium t-butoxide
30 (0.29 ml of a 1 N solution in THF, Aldrich) was added. The mixture was stirred at 125° C overnight. The solvent was evaporated and the residue was purified by flash column chromatography (5 cm; chloroform/ethyl acetate, 2:1) to give the product as a slightly yellowish foam (1.26 g, 80% from
35 starting 2-thiouracil). ¹H-NMR (CDCl₃): δ 13.9 (b m, 1 H,

WO 95/24185

CT/US9: 62652

- 23 -

N··H··O); 7.25 (s, 3 H, CH₃); 4.98 (s, 1 H, CH); 3.5 (s, 3 H, OCH₃); 2.2 (s, 1 H, COCH₃).

Example 12

2-(Hexan-2-on-1-yl)-1-{2'-deoxy-3'-O-[(tert-butyl)diphenylsilyl]-β-D-erythro-furanosyl}-pyrimidin-4-one

2-Hexanone (1.50 ml, 12 mmol) in THF (60 ml) was cooled to -70° C. Lithium diisopropylamide (2 M sol in THF, Aldrich) was added and the solution was stirred for 20 min maintaining the -70° C temperature. The solution was warmed to room temperature and O2,5'-anhydro-3'-O-(tert-butyl)diphenylsilyl)thymidine (1.74 g, 3.76 mmol) was added. The resulting solution became yellowish and after 35 min TLC indicated complete reaction. Water was added and the mixture neutralized carefully with 1 N HCl (alternately 3N acetic acid or 1 N H₂SO₄ can be used). The reaction mixture was partitioned between ether (150 ml) and water (50 ml) and layers separated. The organic layer was dried over Na₂SO₄, filtered and evaporated. The residue was purified by flash column chromatography (5 cm, ethylacetate) to yield the product as a slightly yellowish foam (1.47 g, 69%). ¹H-NMR (CDCl₃): δ 14.1 (s, 1 H, N··H··O); 5.8 (m, 1 H, H-1'); 4.8 (s, 1 H, CH); 2.3 (m, 1 H, CH₂); 1.8 (m, 2 H, OH & CH₃ 5-pyr); 1.4, 1.6 (m, 4 H, CH₂); 0.9 (t, 3 H, CH₃ chain).

Example 13

2-(Hexan-2-on-1-yl)-1-(2'-deoxy-β-D-erythro-furanosyl)-pyrimidin-4-one

2-(Hexan-2-on-1-yl)-1-{2'-deoxy-3'-[(tert-butyl)diphenylsilyl]-β-D-erythro-furanosyl}-pyrimidin-4-one in THF was deblocked with tetrabutylammonium fluoride (Bu₄NF) using standard conditions (see, e.g., Greene and Wuts in *Protective Groups in Organic Synthesis*, 2d ed., John Wiley & Sons, Inc., 1991, p. 83) to give the title compound. ¹H-NMR (CDCl₃): δ 14.3 (s, 1 H, N··H··O); 5.8 (t, 1 H, H1'); 5.35, 5.15 (d,t, 2 H, 3' OH & 5' OH).

WO 95/24185

PCJ 7/S94/02652

- 24 -

Example 14

2-Butyl-1-{2'-deoxy-3'-O-[(tert-butyl)diphenylsilyl]-β-D-erythro-furanosyl}-pyrimidin-4-one

A solution of O-2,5'-anhydro-3'-O-(tert-butyl-diphenyl-silyl)thymidine (0.87 g, 1.88 mmol) in THF (50 ml) was cooled to -70° C and butyllithium (3.75 ml of a 1.6 M solution in hexanes, Aldrich) was added. The mixture turned dark yellow immediately. TLC after 30 mins indicated complete reaction. The reaction mixture was warmed to room temperature, quenched with water (5 ml) and neutralized with 0.1 N HCl. The mixture was extracted with ether, the organic phase washed with H₂O, dried over Na₂SO₄ and evaporated to give a yellow oil (1.08 g). The oil was further purified by flash column chromatography (3 cm, chloroform/methanol 9:1) to give the product as a yellow foam (0.77 g, 79%). ¹H-NMR (CDCl₃): δ 6.1 (t, 1 H, H1'); 2.9 (t, 2 H, CH₂); 2.5 (br m, 1 H, 5' OH); 1.9 (s, 3 H, CH₃ pyr); 1.7, 1.4 (m, 4 H, CH₂); 0.9 (t, 3 H, CH₃ chain).

Example 15

2-Methyl-1-(2',3',5'-tri-O-benzoyl-β-D-ribofuranosyl)-pyrimidin-4-one

2-Methylpyrimidin-4-one (4.42 g, 40 mmol) and a few crystals of ammonium sulfate were refluxed in HMDS overnight. After cooling to room temperature the colorless solution was evaporated. A solution of 1-O-acetyl-2,3,5-tri-O-benzoyl-β-D-ribose (40 mmol, 20.18 g) in 1,2-dichloroethane (200 ml) was added to the residue. Trimethylsilyl trifluoromethanesulfonate (TMSOTf, 9.78 g, 44 mmol) was added and the resulting clear solution was stirred at room temperature for 2 hrs. The reaction mixture was poured in to saturated NaHCO₃ (300 ml) and extracted with chloroform (2 x 200 ml). The organic phase was evaporated to give a colorless homogeneous foam (23.04 g, 104%, pure by NMR).

WO 95/24185

PCT/US94/02652

- 25 -

Example 16**2-Methyl-1-(- β -D-ribofuranosyl)pyrimidin-4-one**

To a solution of 2-methyl-1-(2',3',5'-tri-O-benzoyl- β -D-ribofuranosyl)-pyrimidin-4-one (13.86 g, 25 mmol) in methanol (150 ml) was added 4.37 N sodium methoxide solution in MeOH (2.5 mmol, 0.1 eq.). The solution was stirred overnight at room temperature and then evaporated. The residue was partitioned between water (100 ml) and dichloromethane (2 x 150 ml). The aqueous layer was neutralized with Dowex 50 H⁺-form and evaporated. The orange oily residue was crystallized from ethanol/acetone 3:1 to give the title compound (2.37 g, 39%).
¹H-NMR (CDCl₃): δ 8.1 (d, 1 H, H6); 6.0 (d, 1 H, H5); 5.65 (m, 2 H, H1', 2'-OH); 5.25, 5.2 (d,t, 2 H, 3'-OH, 5'-OH); 2.45 (s, 3 H, CH₃).

Example 17**5-[(tert-Butyl)dimethylsilyl]-2,3-isopropylidene-D-ribo-1,4-lactone**

The title compound was prepared in a manner similar to the preparation of 5-[(tert-butyl)diphenylsilyl]-2,3-isopropylidene-D-ribo-1,4-lactone as reported by Piccirilli, et al., Helvetica Chimica Acta 1991, 74, 397-406, substituting 5-(tert-butyl)diphenylsilyl chloride for the 5-(tert-butyl)diphenylsilyl chloride. The product was obtained as a crystalline compound.

Example 18**3-{5'-O-[(tert-Butyl)dimethylsilyl]-1'-hydroxyl-2',3'-O-isopropylidene-D-ribofuranosyl}-2-chloro-6-methoxy-pyridine and 5-{5'-O-[(tert-Butyl)dimethylsilyl]-1'-hydroxyl-2',3'-O-isopropylidene-D-ribofuranosyl}-2-chloro-6-methoxy-pyridine**

Lithium diisopropylamide (12.6 mmol, 6.3 ml of a 2 M solution) was added to THF (30 ml) at -70° C. To the resulting solution was added 2-chloro-6-methoxypyridine (1.5 g, 1.25 ml, 10.50 mmol) and the resulting orange, clear solution was stirred at -70° C in a dry ice bath for 45 mins during which time the solution slowly turned yellow. 5-[(tert-Butyl)-

WO 95/24185

US 95/02652

- 26 -

dimethylsilyl]-2,3-isopropylidene-D-ribo-1,4-lactone (1.44 g, 5.25 mmol) was added and after 30 mins the dry ice bath was removed and the reaction mixture slowly turned brown as it warmed to room temperature. After 3 hrs at room temperature the reaction was quenched with water and extracted with ether. The combined organic phase was washed with brine, dried over MgSO_4 and evaporated to give a yellow oil (2.28 g) of a mixture of the α and β isomers of the title compounds.

Example 19

3-{5'-O-[(*tert*-Butyl)dimethylsilyl]-2',3'-O-isopropylidene-D-ribofuranosyl}-2-chloro-6-methoxy-pyridine

To a solution of the isomeric mixture of the above 1'-hydroxy compounds in toluene under N_2 in an ice bath will be added triethylsilane (Et_3SiH , 2 equiv.) and $\text{BF}_3 \cdot \text{Et}_2\text{O}$. The mixture is stirred and allowed to come to room temperature. After 2 hours the mixture will be chromatographed on a silica gel column eluted with ethyl acetate/hexanes. The fractions containing the desired β isomer are pooled to yield the product.

Example 20

3-{5'-O-[(*tert*-Butyl)dimethylsilyl]-2',3'-O-isopropylidene- β -D-ribofuranosyl}-2-(1-hexyl-1-yl)-6-methoxy-pyridine

A suspension of bis-triphenylphosphine palladium (II) and copper (I) iodide in THF under argon is stirred for 10 mins. To this suspension will be added 1-hexyne and triethylamine. The resulting suspension will be stirred for an additional 10 mins followed by the addition of 3-{5'-O-[(*tert*-butyl)dimethylsilyl]-2',3'-O-isopropylidene-D-ribofuranosyl}-2-chloro-6-methoxy-pyridine. The resulting mixture is stirred for 1 hr, filtered and evaporated. The residue is dissolved in diethylether, treated with cold 0.1 N HCl and the organic phase is dried over MgSO_4 . Evaporation of the organic phase will yield the title compound.

WO 95/24185

PCT/US94/02653

- 27 -

Example 21

2-(2-Amino-1-hexen-1-yl)-3-{5'-O-[(tert-Butyl)dimethylsilyl]-2',3'-O-isopropylidene-β-D-ribofuranosyl}-6-methoxy-pyridine 3-{5'-O-[(tert-Butyl)dimethylsilyl]-2',3'-O-isopropylidene-D-ribofuranosyl}-2-(1-hexyl-1-yl)-6-methoxy-pyridine will be treated in a pressure bottle with methanolic ammonia (saturated at -20° C) for 18 hours with stirring. The solvent will be evaporated and the residue purified by flash chromatography to give the title compound.

10 Example 22

2-(2-Amino-1-hexen-1-yl)-3-(2',3'-O-isopropylidene-β-D-ribofuranosyl)-6-methoxy-pyridine

2-(2-Amino-1-hexen-1-yl)-3-{5'-O-[(tert-butyl)-dimethylsilyl]-2',3'-O-isopropylidene-D-ribofuranosyl}-6-methoxy-pyridine in THF will be treated with Bu₄NF at room temperature for 30 mins. The resulting solution will be evaporated and the residue chromatographed on silica gel eluted with MeOH/CHCl₃ to give the title compound.

Example 23

20 2-(2-Amino-1-hexen-1-yl)-3-(β-D-ribofuranosyl)-pyrid-6-one
2-(2-Amino-1-hexen-1-yl)-3-(2',3'-O-isopropylidene-β-D-ribofuranosyl)-6-methoxy-pyridine will be dissolved in water and treated with 1 N HCl to remove the isopropylidene groups and concurrently hydrolyze the methoxy function to the corresponding 6-pyridone. The resulting solution is extracted with CH₂Cl₂, the organic phase is washed with saturated bicarbonate, dried over MgSO₄ and evaporated. The residue is purified by chromatography on a silica gel column. The product containing fractions will be pooled and evaporated to yield the title compound.

WO 95/24185

PCT/US/94/02652

- 28 -

Example 24

3,5-[O-(1,1,3,3-Tetraisopropyl)-1,3-disiloxanediyl]-D-ribo-
1,4-lactone

D-Ribono-1,4-lactone and 1,3-dichloro-1,1,3,3-tetra-
5 isopropyldisiloxane are stirred together in dry pyridine for 16
hrs at 25° C. The solvent is evaporated and the resulting oil
will be dissolved in CH₂Cl₂, washed with saturated sodium bicar-
bonate and the organic layer added to a silica gel column. The
product is eluted from the column with methylene chloride/meth-
10 anol and the appropriate fractions, combined and evaporated to
yield the title compound.

Example 25

2-O-Methyl-3,5-[-O-(1,1,3,3-tetraisopropyl)-1,3-disiloxane-
diyl]-D-ribo-1,4-lactone

15 3,5-[O-(1,1,3,3-Tetraisopropyl)-1,3-disiloxanediyl]-D-
ribo-1,4-lactone will be treated with CH₃I and Ag₂O at 40° C
overnight generally in accordance with the procedure of Inoue,
et al., *Nucleic Acids Research* 1987, 15, 6131 to give the title
compound.

20. Example 26

3-{2-O-Methyl-3,5-[-O-(1,1,3,3-tetraisopropyl)-1,3-disiloxane-
diyl]-D-ribofuranosyl}-2-chloro-6-methoxy-pyridine and 5-{2-O-
Methyl-3,5-[-O-(1,1,3,3-tetraisopropyl)-1,3-disiloxanediyl]-D-
ribofuranosyl}-2-chloro-6-methoxy-pyridine

25 Lithium diisopropylamide (as a 2 M solution) is added
to THF (30 ml) at -70° C. To the resulting solution is added
2-chloro-6-methoxypyridine and the resulting solution stirred
at -70° C in a dry ice bath for 45 mins. 2-O-Methyl-3,5-[-O-
(1,1,3,3-tetraisopropyl)-1,3-disiloxanediyl]-D-ribo-1,4-
30 lactone will be added and after 30 mins the dry ice bath
removed and the reaction mixture slowly allowed to warm to room
temperature. After 3 hrs at room temperature the reaction is
quenched with water and extracted with ether. The combined
organic phase is washed with brine and dried over MgSO₄ and

WO 95/24185

PCT/US 95/00612

- 29 -

evaporated to yield an oil that of a mixture of the α and β isomers of the title compounds.

Example 27

3-{2-O-Methyl-3,5-[-O-(1,1,3,3-tetraisopropyl)-1,3-disiloxane-
5 diyl]- β -D-ribofuranosyl}-2-chloro-6-methoxy-pyridine

To a solution of the isomeric mixture of the corresponding 1'-hydroxy compounds in toluene under N_2 in an ice bath is added Et_3SiH (2 equiv.) and $BF_3 \cdot Et_2O$. The mixture is stirred and allowed to come to room temperature. After 2
10 hours the mixture will be chromatographed on a silica gel column eluted with ethyl acetate/hexanes. The fractions containing the desired 3 β isomer are pooled to yield the product.

Example 28

15 3-{2-O-Methyl-3,5-[-O-(1,1,3,3-tetraisopropyl)-1,3-disiloxane-
diyl]- β -D-ribofuranosyl}-2-(1-hexyn-1-yl)-6-methoxy-pyridine

A suspension of bis-triphenylphosphine palladium (II) and copper (I) iodide in THF under argon is stirred for 10 mins. To this suspension will be added 1-hexyne and triethyl-
20 amine. The resulting suspension will be stirred for an additional 10 mins followed by the addition of 3-{2-O-methyl-3,5-[-O-(1,1,3,3-tetraisopropyl)-1,3-disiloxanediyl]-D-ribofuranosyl}-2-chloro-6-methoxy-pyridine. The resulting mixture is stirred for 1 hr, filtered and evaporated. The residue is
25 dissolved in diethylether, treated with cold 0.1 N HCl and the organic phase is dried over $MgSO_4$. Evaporation of the organic phase will yield the title compound.

Example 29

30 3-{2-O-Methyl-3,5-[-O-(1,1,3,3-tetraisopropyl)-1,3-disiloxane-
diyl]- β -D-ribofuranosyl}-6-methoxy-2-(2-thiol-1-hexen-1-yl)-
pyridine

3-{2-O-Methyl-3,5-[-O-(1,1,3,3-tetraisopropyl)-1,3-disiloxanediyl]-D-ribofuranosyl}-2-(1-hexyl-1-yl)-6-methoxy-pyridine in THF will be treated with a steady stream of H_2S for

WO 95/24185

PCT/US 94.02652

- 30 -

3 mins with stirring. The solvent will be evaporated and the residue purified by flash chromatography to give the title compound.

Example 30

5 3-(2-O-Methyl- β -D-ribofuranosyl)-2-(2-thiol-1-hexen-1-yl)-pyrid-6-one

3-{2-O-Methyl-3,5-[-O-(1,1,3,3-tetraisopropyl)-1,3-disiloxanediyl]-D-ribofuranosyl}-6-methoxy-2-(2-thiol-1-hexen-1-yl)-pyridine will be treated with 3 N HCl to remove the
10 tetraisopropylidisiloxanediyl protecting group and hydrolyze the 6-methoxy moiety. The resulting solution will be extracted with CH₂Cl₂, washed with saturated bicarbonate, dried over MgSO₄, and the organic phase evaporated. The residue will be chromatographed on silica gel eluted with MeOH/CHCl₃ to give the title
15 compound.

Those skilled in the art will appreciate that numerous changes and modifications may be made to the preferred embodiments of the invention and that such changes and modifications may be made without departing from the spirit of
20 the invention. It is therefore intended that the appended claims cover all such equivalent variations as fall within the true spirit and scope of the invention.


WO 95/24185

PCT/US 95/02642

- 31 -

What is claimed is:

1. A compound of the structure:



wherein:

R₅ is H or CH₃;

- 5 one of R₂ and R₄ is =O, =NH, or =NH₂⁺; and the other of R₂ and R₄ is Q, =C(R_A)-Q, C(R_A)(R_B)-C(R_C)(R_D)-Q, C(R_A)=C(R_C)-Q or C≡C-Q;

R_A, R_B, R_C and R_D independently are H, SH, OH, NH₂, or C₁-C₂₀ alkyl, or one of (R_A)(R_B) or (R_C)(R_D) is =O;

- 10 Q is halogen, hydrogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkylamine, C₁-C₂₀ alkyl-N-phthalimide, C₁-C₂₀ alkylimidazole, C₁-C₂₀ alkylbis-imidazole, imidazole, bis-imidazole, amine, N-phthalimide, C₂-C₂₀ alkenyl, C₂-C₂₀ alkynyl, hydroxyl, thiol, keto, carboxyl, nitrates, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, O-(hydroxyl protecting group), a leaving group, a heterocycle, an intercalator, a reporter molecule, a conjugate, a polyamine, a polyamide, a polyethylene glycol, a polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, a group that enhances the pharmacokinetic properties of oligonucleotides, a RNA cleaving moiety or a depurination enhancing group;
- 20

WO 95/24185

PCT/US94/(2)

- 32 -

T_3 and T_5 independently are H, phosphate, an activated phosphate, a hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleotide or an oligonucleoside;

X_1 and X_2 independently are hydrogen, halogen, hydroxyl, O-(hydroxyl protecting group), thiol, carboxyl, nitrate, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocyclic, alicyclic, carbocyclic, intercalators, reporter molecules, conjugates, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligonucleotides, or groups that enhance the pharmacokinetic properties of oligonucleotides; and

provided that when T_5 is a hydroxyl protecting group or together T_3 and one of X_1 or X_2 is a hydroxyl protecting group, then R_2 is not CH_3 ; and that when T_3 and T_5 are OH or O-benzoyl, then R_2 is not $CH_2OC_6H_5$; and that when R_4 is Q, then Q is not hydrogen, hydroxyl, thiol, amino or hydrogen.

2. A compound of claim 1 wherein one of X_1 and X_2 is hydrogen and the other is hydroxyl.

3. A compound of claim 1 wherein both of X_1 and X_2 are hydrogen.

4. A compound of claim 1 wherein R_2 is halogen.

5. A compound of claim 4 wherein R_2 is chlorine.

6. A compound of claim 1 wherein R_4 is =O.

7. A compound of claim 1 wherein R_5 is H.

8. A compound of claim 1 wherein R_5 is CH_3 .

WO 95/24183

PCT/US94/02652

- 33 -

9. A compound of claim 1 wherein T_1 and T_2 are both H.

10. A compound of claim 1 wherein T_1 is an activated phosphate and T_2 is a hydroxyl protecting group.

5 11. A compound of claim 1 wherein at least one of T_1 or T_2 is an oligonucleotide or an oligonucleoside.

12. A compound of claim 1 wherein one of R_2 or R_4 is $C(R_A)(R_B)-C(R_C)(R_D)-Q$.

10 13. A compound of claim 12 wherein R_A and R_B are both hydrogen, and $(R_C)(R_D)$ is $=O$.

14. A compound of claim 13 wherein Q includes a sulfur containing heterocycle and a linking group covalently bound between said C(O) and said sulfur containing heterocycle.

15 15. A compound of claim 13 wherein said linking group is C_1-C_6 alkyl, C_2-C_{20} alkenyl or C_2-C_{20} alkynyl.

16. A compound of claim 1 wherein one of R_2 or R_4 is $C(R_A)=C(R_C)-Q$.

17. A compound of claim 16 wherein R_A is OH, SH or NH_2 ; and R_C is hydrogen.

20 18. A compound of claim 17 wherein Q is C_1-C_{20} alkyl.


19. A compound of claim 17 wherein Q is C_1-C_6 alkyl.

WO 95/24185

PCT/US94/02652

- 34 -

20. A compound of the structure:



wherein:

R_5 is H or CH_3 ;

one of R_2 and R_4 is $=\text{O}$, $=\text{NH}$, or $=\text{NH}_2$; and the other of

5 R_2 and R_4 is $=\text{C}(\text{R}_\text{A})-\text{Q}$, $\text{C}(\text{R}_\text{A})(\text{R}_\text{B})-\text{C}(\text{R}_\text{C})(\text{R}_\text{D})-\text{Q}$ or $\text{C}(\text{R}_\text{A})=\text{C}(\text{R}_\text{C})-\text{Q}$;

R_A , R_B , R_C and R_D independently are H, SH, OH or NH_2 ,
or one of $(\text{R}_\text{A})(\text{R}_\text{B})$ or $(\text{R}_\text{C})(\text{R}_\text{D})$ is $=\text{O}$;

Q is halogen, hydrogen, $\text{C}_1\text{-C}_{20}$ alkyl, $\text{C}_1\text{-C}_{20}$ alkylamine,
 $\text{C}_1\text{-C}_{20}$ alkyl-N-phthalimide, $\text{C}_1\text{-C}_{20}$ alkylimidazole, $\text{C}_1\text{-C}_{20}$
10 alkylbis-imidazole, imidazole, bis-imidazole, amine or N-phthalimide;

T_3 and T_5 independently are H, phosphate, an activated
phosphate, hydroxyl protecting group, a nucleoside, a
nucleotide, an oligonucleotide or an oligonucleoside;

15 X_1 and X_2 independently are hydrogen, hydroxyl or O-
(hydroxyl protecting group); and

provided that when T_5 is a hydroxyl protecting group
or together T_3 and one of X_1 or X_2 is a hydroxyl protecting
group, then R_2 is not CH_3 ; and that when T_3 and T_5 are OH or O-
20 benzoyl, then R_2 is not $\text{CH}_2\text{OC}_6\text{H}_5$ and the when R_4 is Q, Q is not
hydrogen, hydroxyl, thiol, amino or hydrogen..

21. A compound of claim 20 wherein one of X_1 and X_2
is hydrogen and the other is hydroxyl.

22. A compound of claim 20 wherein both of X_1 and X_2
25 are hydrogen.

WO 95/24185

PCT/US94/02652

- 35 -


23. A compound of claim 20 wherein R_2 is halogen.
24. A compound of claim 23 wherein R_2 is chlorine.
25. A compound of claim 20 wherein R_4 is =O.
26. A compound of claim 20 wherein R_5 is H.
- 5 27. A compound of claim 20 wherein R_5 is CH_3 .
28. A compound of claim 20 wherein T_3 and T_5 are both H.
29. A compound of claim 20 wherein T_3 is an activated phosphate and T_5 is a hydroxyl protecting group.
- 10 30. A compound of claim 20 wherein at least one of T_3 or T_5 is an oligonucleotide or an oligonucleoside.
31. A compound of claim 20 wherein one of R_2 or R_4 is $C(R_A)(R_B)-C(R_C)(R_D)-Q$.
- 15 32. A compound of claim 31 wherein R_A and R_B are both hydrogen; and $(R_C)(R_D)$ is =O.
33. A compound of claim 20 wherein one of R_2 is or R_4 is $C(R_A)=C(R_C)-Q$.
34. A compound of claim 33 wherein R_A is OH, SH or NH_2 ; and R_C is hydrogen.

WO 95/14185

PCT/US94/02652

- 36 -

35. An oligonucleoside containing at least one nucleoside of the structure:



wherein:

- R_5 is H or CH_3 ;
- 5 one of R_2 and R_4 is $=\text{O}$, $=\text{NH}$, or $=\text{NH}_2^+$; and the other of R_2 and R_4 is Q, $\text{C}(R_A)(R_B)-\text{C}(R_C)(R_D)-\text{Q}$, $\text{C}(R_A)=\text{C}(R_C)-\text{Q}$ or $\text{C}\equiv\text{C}-\text{Q}$;
- R_A , R_B , R_C and R_D independently are H, SH, OH, NH_2 or C_1-C_{20} alkyl, or one of $(R_A)(R_B)$ or $(R_C)(R_D)$ is $=\text{O}$;
- Q is halogen, hydrogen, C_1-C_{20} alkyl, C_1-C_{20} alkylamine,
- 10 C_1-C_{20} alkyl-N-phthalimide, C_1-C_{20} alkylimidazole, C_1-C_{20} alkylbis-imidazole, imidazole, bis-imidazole, N-phthalimide, C_2-C_{20} alkenyl, C_2-C_{20} alkynyl, hydroxyl, thiol, keto, carboxyl, nitrates, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-
- 15 aralkyl, NH-aralkyl, amino, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, O-(hydroxyl protecting group), a leaving group, a heterocycle, an intercalator, a reporter molecule, a conjugate, a polyamine, a polyamide, a polyethylene glycol, a polyether, a group that
- 20 enhance the pharmacodynamic properties of oligonucleotides, a group that enhance the pharmacokinetic properties of oligonucleotides, a RNA cleaving moiety or a depurination enhancing group;
- X_1 and X_2 independently are hydrogen, halogen, hydrox-
- 25 yl, O-(hydroxyl protecting group), thiol, carboxyl, nitrate, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-

WO 95/24185


PCT/US94/02652

- 37 -

aralkyl, amino, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocyclic, alicyclic, carbocyclic, intercalators, reporter molecules, conjugates, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligonucleotides, groups that enhance the pharmacokinetic properties of oligonucleotides; and

T_3 and T_5 independently are H, a nucleoside or an oligonucleoside.

- 10 36. An oligonucleoside containing at least one nucleoside of the structure:



wherein:

R_5 is H or CH_3 ;

one of R_2 and R_4 is $=O$, $=NH$, or $=NH_2^+$; and the other of

- 15 R_2 and R_4 is $C(R_A)(R_B)-C(R_C)(R_D)-Q$ or $C(R_A)=C(R_C)-Q$;

R_A , R_B , R_C and R_D independently are H, SH, OH or NH_2 , or one $(R_A)(R_B)$ or $(R_C)(R_D)$ is $=O$;

- Q is halogen, hydrogen, C_1-C_{20} alkyl, C_1-C_{20} alkylamine, C_1-C_{20} alkyl-N-phthalimide, C_1-C_{20} alkylimidazole, C_1-C_{20} alkylbis-imidazole, imidazole, bis-imidazole, amine or N-phthalimide;

X_1 and X_2 independently are hydrogen, hydroxyl or O-(hydroxyl protecting group); and

- 25 T_3 and T_5 independently are H, a nucleoside, or an oligonucleoside that is joined to said structure.

WO 95/24185

PCT/US94/02652


- 38 -

37. A process of treating an organism having a disease characterized by the undesired production of a protein, comprising contacting the organism with a compound of claim 1.

38. A method of modulating the production or activity of a protein in an organism comprising contacting the organism with a compound of claim 1.

39. A pharmaceutical composition comprising a compound of claim 1 in a pharmaceutically effective carrier.

40. A compound of the structure:



10 wherein:

J is N or CH;

R₄ is =O, =NH, or =NH₂⁺;

R₂ is Q, =C(R_A)-Q, C(R_A)(R_B)-C(R_C)(R_D)-Q, C(R_A)=C(R_C)-Q or C≡C-Q;

15 R_A, R_B, R_C and R_D independently are H, SH, OH, NH₂ or C₁-C₂₀ alkyl, or one of (R_A)(R_B) or (R_C)(R_D) is =O;

Q is halogen, hydrogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkylamine, C₁-C₂₀ alkyl-N-phthalimide, C₁-C₂₀ alkylimidazole, C₁-C₂₀ alkylbis-imidazole, imidazole, bis-imidazole, amine, N-phthalimide, C₂-C₂₀ alkenyl, C₂-C₂₀ alkynyl, hydroxyl, thiol, keto, carboxyl, nitrates, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide,

WO 95/24185

PCT/US94/02652

- 39 -

disulfide, silyl, O-(hydroxyl protecting group), a leaving group, a heterocycle, an intercalator, a reporter molecule, a conjugate, a polyamine, a polyamide, a polyethylene glycol, a polyether, a group that enhance the pharmacodynamic properties of oligonucleotides, a group that enhance the pharmacokinetic properties of oligonucleotides, a RNA cleaving moiety or a depurination enhancing group;

T₃ and T₅ independently are H, phosphate, an activated phosphate, hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleotide or an oligonucleoside; and

X₁ and X₂ independently are hydrogen, halogen, hydroxyl, O-(hydroxyl protecting group), thiol, carboxyl, nitrate, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocyclic, alicyclic, carbocyclic, intercalators, reporter molecules, conjugates, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligonucleotides, or groups that enhance the pharmacokinetic properties of oligonucleotides.

41. A method of preparing a compound according to claim 1, comprising contacting a 2,2'-anhydropyrimidine with a nucleophile having formula R₂ under conditions effective to form a covalent bond between said nucleophile and the 2-position of said pyrimidine.

42. A method of preparing a compound according to claim 1, comprising contacting a 2,5'-anhydropyrimidine with a nucleophile having formula R₂ under conditions effective to form a covalent bond between said nucleophile and the 2-position of said pyrimidine.

43. A method of preparing a compound according to claim 1 wherein R₁ is C≡C-Q, comprising contacting a 4-halopyrimidine with an alkyne having formula HC≡C-Q in the presence

WO 95/24185

P 71/US94/12652

- 40 -

of an organometallic coupling reagent for a time and under reaction conditions effective to form a covalent bond between said alkyne and the 2-position of said pyrimidine.

WO 95/24185

FCT/US 4/02652

1/1




FIGURE 1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/02652

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :A61K 31/00, 48/00

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44, 49, 50,51; 536/4.1,22.1,25.3, 25.31, 25.33, 26.8, 28.1, 28.5, 28.51

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS on line

APS on line

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 3,755,295 (VERHEYDEN ET AL) 28 August 1973, see columns 1-14.	1-34,39-42
Y	US, A, 3,975,374 (HUNTER ET AL) 17 AUGUST 1976, see columns 1,2,6,7,9,10,14, 26-28.	1-34,37,39-42
Y	US, A, 5,212,295 (COOK ET AL) 18 May 1993, see columns 1-32.	1-43



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	Z	document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

15 JULY 1994

Date of mailing of the international search report

AUG 01 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

JAMES O. WILSON

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/02652

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

514/44, 49, 50, 51; 536/4.1, 22.1, 25.3, 25.31, 25.33, 26.8, 28.1, 28.5, 28.51